

Updated Analysis of a_2/a_1 in Exclusive B Decays ^a

Hai-Yang Cheng and Kwei-Chou Yang

Institute of Physics, Academia Sinica, Taipei, Taiwan 115, R.O.C.

Using recent experimental data and various theoretical calculations on form factors, we reanalyze the effective parameters a_1 and a_2 and their ratio.

Nonleptonic two-body decays of mesons were conventionally studied in the generalized factorization approach, in which the factorized decay amplitude is associated with the effective coefficients

$$a_{1,2}^{\text{eff}} = c_{1,2}^{\text{eff}} + c_{2,1}^{\text{eff}}[(1/N_c) + \chi_{1,2}], \quad (1)$$

where $c_{1,2}^{\text{eff}}$ are the effective Wilson coefficients and the nonfactorized terms are expressed by $\chi_{1,2}$. For simplicity, in what follows we will drop the superscript “eff” of $a_{1,2}^{\text{eff}}$. Based on the generalized factorization assumption, one can catalog the decay processes into three different classes. For class-I decays, the decay amplitudes, dominated by color-allowed external W-emission, are proportional to the a_1 parameter. For class-II decays, the decay amplitudes, governed by color-suppressed internal W-emission, are described by a_2 . The decay amplitudes of the class-III involve a linear combination of a_1 and a_2 . From the data analysis, contrary to the case of charmed meson decays in which a_2 is negative and $\chi \sim -1/N_c$, a_2 is positive in two-body decays of the B meson. $\chi_{2(1)}$ coming from the contribution of the matrix element of the color octet-octet and singlet-singlet currents, since it is made of nonfactorizable soft gluon effects in itself, will become more important if the final particles move slower^{1,2}. Therefore, we naïvely expect that $|\chi_i(D \rightarrow V P)| \gtrsim |\chi_i(D \rightarrow P P)| > |\chi_i(B \rightarrow V P)|$, where P is the pseudoscalar meson and V the vector meson. The data of $B \rightarrow J/\Psi K^{(*)}$ which are class-II decay modes can be used to study the absolute value of a_2 . However, the experimental results are quite difficult to account for the observed longitudinal polarization fraction Γ_L/Γ and the ratio $R \equiv \Gamma(B \rightarrow J/\Psi K^*)/\Gamma(B \rightarrow J/\Psi K)$ simultaneously

^apresented by K.-C. Yang at “The Fourth International Workshop on Particle Physics Phenomenology”, Kaohsiung, Taiwan, 18-21 January 1998.

Email address: kcyang@phys.sinica.edu.tw

Table 1: The effective parameter a_1 extracted from $B \rightarrow D^{(*)} D_s^{(*)}$ using different form-factor models.

	BSWI	BSWII	LF	SR+HQET	NS
$\bar{B}^0 \rightarrow D^+ D_s^-$	1.05 ± 0.16	1.05 ± 0.16	1.06 ± 0.16	1.17 ± 0.18	1.32 ± 0.20
$B^- \rightarrow D^0 D_s^-$	0.80 ± 0.15	0.80 ± 0.15	0.81 ± 0.15	0.89 ± 0.17	1.01 ± 0.19
$\bar{B}^0 \rightarrow D^+ D_s^{*-}$	1.24 ± 0.31	1.10 ± 0.27	1.11 ± 0.28	1.26 ± 0.31	1.43 ± 0.36
$B^- \rightarrow D^0 D_s^{*-}$	1.14 ± 0.25	1.10 ± 0.22	1.02 ± 0.23	1.16 ± 0.26	1.32 ± 0.29
$\bar{B}^0 \rightarrow D^{*+} D_s^-$	1.34 ± 0.24	1.21 ± 0.21	1.11 ± 0.28	1.09 ± 0.19	1.03 ± 0.18
$B^- \rightarrow D^{*0} D_s^-$	1.46 ± 0.30	1.31 ± 0.28	1.02 ± 0.23	1.18 ± 0.25	1.12 ± 0.23
$\bar{B}^0 \rightarrow D^{*+} D_s^{*-}$	0.88 ± 0.15	0.90 ± 0.16	0.83 ± 0.15	0.86 ± 0.15	0.85 ± 0.15
$B^- \rightarrow D^{*0} D_s^{*-}$	1.05 ± 0.20	1.07 ± 0.20	0.99 ± 0.18	1.03 ± 0.19	1.02 ± 0.19
Average	1.03 ± 0.07	1.01 ± 0.07	0.97 ± 0.07	1.03 ± 0.07	1.06 ± 0.07

using the existing form factor calculations. To accommodate the data, it has been advocated² a modification to the generalized factorization hypothesis so that a_2 is process dependent. In this talk, we focus on the extraction of the value a_1 , $|a_2|$, and a_2/a_1 from the current experimental data³ and various form factor calculations⁴, the BSWI model, the BSWII model⁵, the LF model⁶, the NS model⁷, the QCD sum rule results (Yang)⁸, and the light-cone sum rule calculations (LC)⁹. Since the accuracy of these values still suffers from reliability of the form factor calculations, we desire that an objective estimation can be obtained.

Here will consider the decay amplitudes of processes which have been measured experimentally:

- Class I: $B^- \rightarrow D^{(*)0} D_s^{(*)-}$, $\bar{B}_d^0 \rightarrow D^{(*)+} D_s^{(*)-}$, and $\bar{B}_d^0 \rightarrow D^{(*)+} \pi^- (\rho^-)$,
- Class II: $B^- \rightarrow J/\Psi K^{(*)-}$ and $B^0 \rightarrow J/\Psi K^{(*)0}$,
- Class III: $B^- \rightarrow D^{(*)0} \pi^- (\rho^-)$,

The value of $a_{(1,2)}$ can be extracted from the above data of the class-(I,II) type transitions, while their ratio can be obtained by the the following ratios:

$$\begin{aligned}
 R_1 &\equiv \frac{\mathcal{B}(B^- \rightarrow D^0 \pi^-)}{\mathcal{B}(\bar{B}^0 \rightarrow D^+ \pi^-)}, & R_2 &\equiv \frac{\mathcal{B}(B^- \rightarrow D^0 \rho^-)}{\mathcal{B}(\bar{B}^0 \rightarrow D^+ \rho^-)}, \\
 R_3 &\equiv \frac{\mathcal{B}(B^- \rightarrow D^{*0} \pi^-)}{\mathcal{B}(\bar{B}^0 \rightarrow D^{*+} \pi^-)}, & R_4 &\equiv \frac{\mathcal{B}(B^- \rightarrow D^{*0} \rho^-)}{\mathcal{B}(\bar{B}^0 \rightarrow D^{*+} \rho^-)}.
 \end{aligned} \tag{2}$$

The hadronic matrix element of effective operators $O_{1,2}$ can be redefined in the scheme- and scale(μ)-dependent ways $\langle O(\mu) \rangle = g(\mu) \langle O \rangle_{\text{tree}}$, where $\langle O \rangle_{\text{tree}}$

Table 2: The effective parameter a_1 extracted from $\bar{B}_d^0 \rightarrow D^{(*)+} \pi^-(\rho^-)$ using different form-factor models.

	BSWI	BWSII	LF	SR+HQET	NS
$\bar{B}^0 \rightarrow D^+ \pi^-$	0.89 ± 0.06	0.89 ± 0.06	0.88 ± 0.06	1.15 ± 0.08	1.31 ± 0.09
$\bar{B}^0 \rightarrow D^+ \rho^-$	0.91 ± 0.08	0.90 ± 0.08	0.89 ± 0.08	1.13 ± 0.10	1.29 ± 0.11
$\bar{B}^0 \rightarrow D^{*+} \pi^-$	0.98 ± 0.04	0.98 ± 0.04	0.84 ± 0.03	0.93 ± 0.04	0.89 ± 0.03
$\bar{B}^0 \rightarrow D^{*+} \rho^-$	0.86 ± 0.21	0.87 ± 0.21	0.75 ± 0.18	0.83 ± 0.20	0.80 ± 0.20
Average	0.95 ± 0.03	0.94 ± 0.03	0.85 ± 0.03	0.98 ± 0.03	0.96 ± 0.03

Table 3: The effective parameter $|a_2|$ extracted from $B \rightarrow J/\Psi K^{(*)}$ using different form-factor models.

	BSWI	BWSII	LF	NS	Yang	LC
$B^+ \rightarrow J/\Psi K^+$	0.33 ± 0.02	0.22 ± 0.01	0.28 ± 0.01	0.36 ± 0.02	0.37 ± 0.02	0.30 ± 0.02
$B^0 \rightarrow J/\Psi K^0$	0.32 ± 0.02	0.21 ± 0.01	0.27 ± 0.02	0.35 ± 0.02	0.36 ± 0.02	0.29 ± 0.02
Average	0.32 ± 0.01	0.22 ± 0.01	0.28 ± 0.01	0.36 ± 0.01	0.36 ± 0.01	0.30 ± 0.01
$B^+ \rightarrow J/\Psi K^{*+}$	0.20 ± 0.02	0.21 ± 0.02	0.25 ± 0.02	0.24 ± 0.02	0.39 ± 0.04	0.20 ± 0.02
$B^0 \rightarrow J/\Psi K^{*0}$	0.19 ± 0.01	0.21 ± 0.01	0.25 ± 0.02	0.25 ± 0.02	0.39 ± 0.03	0.19 ± 0.01
Average	0.20 ± 0.01	0.22 ± 0.01	0.26 ± 0.01	0.25 ± 0.01	0.40 ± 0.02	0.20 ± 0.01

is μ -independent. We thus can rewrite the matrix element of the effective Hamiltonian $\langle H_{\text{eff}} \rangle$ as the product of c^{eff} and $\langle O \rangle_{\text{tree}}$, both of which are scheme- and μ -independent. The only relevant scale, which is implicit and of order m_b , separating c^{eff} and $\langle O \rangle_{\text{tree}}$ is the energy release of B decays. Here we use $c_1^{\text{eff}} = 1.149$ and $c_2^{\text{eff}} = -0.325$, to the next-to-leading order¹⁰. Using the theoretical results of form factors and the experimental data from the Particle Data Group (PDG)³, one can easily extract the values of a_1 , $|a_2|$, and a_2/a_1 . The results are listed in Tables 1, 2, 3, and 5. In Tables 1 and 2, the updated data are used for the parameters of NS, while the results of SR+HQET^b are gotten by the NS formula but the parameters are obtained from the sum rule results and the predictions of the heavy quark effective theory¹¹. All the results^c in Tables 1 and 2 indicate that the value of a_1 , which is very close to 1, seems to follow the pattern $a_1(B \rightarrow D^{(*)} D_s^{(*)}) \gtrsim a_1(B \rightarrow D^{(*)} \pi(\rho))$.

For $B \rightarrow J/\Psi K^{(*)}$, since they are color suppressed and sensitive to the nonfactorizable contributions, therefore, they are very good examples to ex-

^bWe will use the $B \rightarrow D^{(*)}$ form-factor results of SR+HQET as subsidiaries of the Yang's⁸ and LC⁹ calculations, since Yang's sum rules and LC only show the form factors of the B meson into a light meson.

^cThe class-I transitions determine the absolute value of a_1 . However, since this kind of processes is color allowed, we expect the sign of a_1 is the same as that of c_1^{eff} and is therefore positive.

Table 4: R , Γ_L/Γ , and $|P|^2$ from various form factor calculations based on the generalized factorization hypothesis together with the CLEO new data.

	BSWI	BSWII	LF	NS	Yang	LC	CLEO
R	4.15	1.58	1.79	3.17	1.30	3.48	1.45 ± 0.26
Γ_L/Γ	0.57	0.36	0.53	0.49	0.42	0.47	0.52 ± 0.08
$ P ^2$	0.09	0.24	0.09	0.12	0.19	0.23	0.16 ± 0.09

Table 5: a_2/a_1 extracted from the PDG data.

	BSWI	BSWII	LF	NS	Yang	LC
R_1	0.29 ± 0.10	0.29 ± 0.10	0.38 ± 0.13	0.28 ± 0.10	0.29 ± 0.10	0.27 ± 0.10
R_2	0.59 ± 0.30	0.52 ± 0.27	0.56 ± 0.28	0.43 ± 0.22	1.21 ± 0.62	0.34 ± 0.17
R_3	0.23 ± 0.06	0.20 ± 0.06	0.31 ± 0.09	0.30 ± 0.09	0.28 ± 0.08	0.25 ± 0.07
R_4	0.55 ± 0.40	0.67 ± 0.49	0.85 ± 0.62	0.79 ± 0.58	1.50 ± 1.09	0.62 ± 0.45
Average	0.26 ± 0.05	0.23 ± 0.05	0.35 ± 0.07	0.31 ± 0.06	0.30 ± 0.06	0.27 ± 0.05

plore the generalized factorization hypothesis to see if a_2 is universal. On the other hand, since the energy release in $B \rightarrow J/\Psi K$ is close to that in $B \rightarrow J/\Psi K^*$, we expect that the extracted values of $|a_2|$ from these two kinds of processes can be slightly different (the so-called minimal modified factorization hypothesis). However, from Table 3, we know that only BSWII, LF, and Yang's sum rules can satisfy our expectation.

The other ways to examine the generalized factorization hypothesis and/or the quality of various form factor results are to study the production ratio $R = \mathcal{B}(B \rightarrow J/\Psi K^*)/\mathcal{B}(B \rightarrow J/\Psi K)$, the fraction of longitudinal polarization Γ_L/Γ , and the P -wave transverse polarization $|P|^2$ measured in the transversity basis in $B \rightarrow J/\Psi K^*$ decays. The predictions from various form factor calculations based on the generalized factorization hypothesis together with the CLEO new data¹² are shown in Table 4. We find again that only BSWII, LF, and Yang's sum rules can accommodate the data, expect that the value of Γ_L/Γ in BSWII is a little smaller than the CLEO data. Furthermore, if allowing the nonfactorizable contribution χ_2 to be slightly different in $B \rightarrow J/\Psi K^{(*)}$ and in $B \rightarrow D^{(*)} \pi(\rho)$, we find that the values of $a_2(B \rightarrow J/\Psi K^{(*)})$ of BSWII/LF are consistent with that obtained from the analyses of R_{1-4} (see Tables 5 and 6) if assuming $a_1 \approx 1$. However, note that the Yang's sum rule results do not behave like the BSWII or LF model. The value of a_2 in the sum rule analysis of Yang is larger. There are two possibilities to explain it. One is $\chi_2 \approx 0.3$, a larger value compared to that in BSWII or LF ($\chi_2 \approx 0.14$ in BWSII, 0.18 in LF if we have adopted $a_2 > 0$). The other possibility is both the nonfactorizable values χ_2 and a_2 become negative in $B \rightarrow J/\Psi K^{(*)}$, as the D meson decays. The QCD sum rule calculation on the

Table 6: a_2/a_1 , if using the CLEO II data.

	BSWI	BSWII	LF	NS	Yang	LC
R_1	0.33 ± 0.13	0.33 ± 0.13	0.43 ± 0.16	0.32 ± 0.12	0.33 ± 0.13	0.31 ± 0.12
R_2	0.10 ± 0.24	0.09 ± 0.21	0.09 ± 0.23	0.07 ± 0.18	0.20 ± 0.49	0.06 ± 0.14
R_3	0.16 ± 0.08	0.14 ± 0.07	0.22 ± 0.10	0.22 ± 0.10	0.20 ± 0.09	0.18 ± 0.08
R_4	0.33 ± 0.14	0.40 ± 0.18	0.51 ± 0.22	0.47 ± 0.21	0.90 ± 0.39	0.37 ± 0.16
Average	0.22 ± 0.06	0.20 ± 0.05	0.29 ± 0.08	0.25 ± 0.07	0.27 ± 0.07	0.21 ± 0.06

nonfactorizable contribution in $B \rightarrow J/\Psi K$ was reported to be negative¹³, contrary to the generalized nonfactorization hypothesis. However, it is difficult to understand why a_2 is negative in $B \rightarrow J/\Psi K^{(*)}$ while it becomes positive in $B^- \rightarrow D^{(*)} \pi^-(\rho^-)$. One way to solve directly this problem is to evaluate the class-III or class-II decay channels of $B \rightarrow D \pi(\rho)$, within framework of the QCD sum rules to see if χ_2 becomes positive.

From the analyses of R_{1-4} defined in Eq. (2) one can obtain a_2/a_1 . Since we have adopted that a_1 is positive, we can thus determine the sign of a_2 . The results are listed in Table 5 if using PDG data. All of the results show that a_2 is positive. Two remarks are in order. First, a_2/a_1 of $R_{1,3}$ for all of form factor results are quantitatively stable, while that of $R_{2,4}$ have large central values and errors. Second, the prediction of a_2/a_1 in LF is slightly large is because a_1 in LF is a smaller value (see Tables 1 and 2).

CLEO II¹⁴ has also reported the data on various class-III and class-I branching ratios of $B \rightarrow D^{(*)} \pi(\rho)$. We show that the analysis results of a_2/a_1 in Table 6. But now the values of a_2/a_1 from $R_{2,4}$ is much smaller the corresponding values in Table 5. Eventually, if we account of the fact that a_1 is little small in LF, then a_2 is lying in $0.2 - 0.3$, which is a reasonable result for all of form factor calculations.

To conclude, we have used the current experimental data and various theoretical results of form factors to analyze the effective coefficients a_1 and a_2 . Our results have shown that if allowing a minimal modification to the generalized factorization hypothesis, *i.e.*, the nonfactorizable contribution $\chi_{1,2}$ can be slightly different in different decay processes, then we have the following conclusions: (1) $a_1 \approx 1$ and $a_1(B \rightarrow D^{(*)} D_s^{(*)}) \gtrsim a_1(B \rightarrow D^{(*)} \pi(\rho))$. (2) Only the BSW II model, the light-front model, and Yang's sum rules can satisfy this factorization assumption in $B \rightarrow J/\Psi K^{(*)}$ analyses. The results are shown in Tables 3 and 4. (3) From the results of a_2/a_1 listed in Tables 5 and 6, we obtain that a_2 is lying in $0.2 - 0.3$, which is a reasonable result for all of form factor calculations in the $B \rightarrow D^{(*)} \pi(\rho)$ decay processes.

This work was supported in part by the National Science Council of R.O.C.

under Grant No. NSC88-2112-M-001-006.

1. H.Y. Cheng and B. Tseng, *Phys. Rev. D* **51**, 6529 (1995); H.Y. Cheng, *Z. Phys. C* **69**, 647 (1996).
2. H.Y. Cheng, *Phys. Lett. B* **395**, 345 (1997).
3. C. Cao *et al.* (Particle Data Group), *Eur. Phys. J.* **C3**, 1 (1998).
4. For a recent similar work, see F.M. Al-Shamali and A.N. Kamal, hep-ph/9806270.
5. M. Bauer, B. Stech, and M. Wirbel, *Z. Phys. C* **34**, 103 (1987).
6. H.Y. Cheng, C.Y. Cheung, and C.W. Hwang, *Phys. Rev. D* **55**, 1559 (1997).
7. M. Neubert, B. Stech, CERN-TH-97-099, in second edition of Heavy Flavours, ed. by A.J. Buras and M. Lindner, World Scientific, Singapore.
8. The detailed results will be published in a separate work. The basic formulas can be found in K.C. Yang, *Phys. Rev. D* **57**, 2983 (1998).
9. P. Ball and V.M. Braun, CERN-TH/98-162 [hep-ph/9805422].
10. H.Y. Cheng and B. Tseng, *Phys. Rev. D* **58**, 094005 (1998).
11. M. Neubert, CERN-TH-97-024 [hep-ph/9702375]; *Phys. Rep.* **245**, 259 (1994).
12. CLEO Collab, C.P. Jessop *et al.*, *Phys. Rev. Lett.* **79**, 4533 (1997).
13. A. Khodjamirian and R. Rückl, WUE-ITP-97-049 [hep-ph/9801443].
14. J.L. Rodriguez, hep-ex/9801028.